





How to give the right amount of ACETAMINOPHEN (also known as Tylenol) is different depending on which medicine you plan to give.

Dose: For products labeled for children and infants, **DO NOT GIVE MORE THAN 5 DOSES IN 24 HOURS**. For children over 12 years, extra strength tablets/gelcaps can be given every 6 hours. **DO NOT GIVE MORE THAN 6 EXTRA STRENGTH TABLETS/GELCAPS IN 24 HOURS**. Acetaminophen should not be given to children under two years of age without the guidance of a doctor.

Weight →	6 to 11 pounds (3 to 5 kilograms)	12 to 17 pounds (5 to 7 kilograms)	18 to 23 pounds (8 to 10 kilograms)	24 to 35 pounds (11 to 15 kilograms)	36 to 47 pounds (16 to 21 kilograms)	48 to 59 pounds (22 to 26 kilograms)	60 to 71 pounds (27 to 32 kilograms)	72 to 95 pounds (33 to 43 kilograms)	96 pounds or more (44 kilograms or more)
Age →	0 to 3 months	4 to 11 months	12 to 23 months	2 to 3 years	4 to 5 years	6 to 8 years	9 to 10 years	11 years	12 years or older
Infants' Acetaminophen (160 mg / 5 mL)	1.25 mL 	2.5 mL 	3.75 mL 	5 mL 	_____	_____	_____	_____	_____
Children's Acetaminophen (160 mg / 5 mL)	_____	_____	_____	5 mL 	7.5 mL 	10 mL 	12.5 mL 	15 mL 	_____
Children's Acetaminophen Chewables (160 mg)	_____	_____	_____	1 tablet 	1 ½ tablets 	2 tablets 	2 ½ tablets 	3 tablets 	_____
Children's Acetaminophen Dissolvable Packets (160 mg / powder pack)	_____	_____	_____	_____	_____	2 packets 	2 packets 	3 packets 	_____
Adults' Acetaminophen Tablets (325 mg; regular strength)	_____	_____	_____	_____	_____	1 tablet 	1 tablet 	1 tablet 	2 tablets 
Adults' Acetaminophen Tablets (500 mg; extra strength)	_____	_____	_____	_____	_____	_____	_____	_____	2 tablets 

For more information, visit [healthychildren.org/feverpain](https://www.healthychildren.org/feverpain)







healthychildren.org

Powered by pediatricians. Trusted by parents.
from the American Academy of Pediatrics

How to give the right amount of IBUPROFEN (also known as Motrin, Advil) is different depending on which type of ibuprofen you plan to give.

Dose: Give every 6 hours if needed, for fever or pain. **DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.**
Do NOT use with any other medicine containing ibuprofen.

Weight	Age	Infant's Ibuprofen Drops (50 mg / 1.25 mL)	Children's Liquid Ibuprofen (100 mg / 5 mL)	Children's Ibuprofen Chewable Tablets (100 mg)	Adult's Ibuprofen Tablets (200 mg)
0 to 11 pounds (up to 5 kilograms)	0 to 5 months	_____	_____	_____	_____
12 to 17 pounds (about 6 to 7 kilograms)	6 to 11 months	1.25 mL 	2.5 mL* 	_____	_____
18 to 23 pounds (about 8 to 10 kilograms)	12 to 23 months	1.875 mL 	4 mL* 	_____	_____
24 to 35 pounds (about 11 to 15 kilograms)	2 to 3 years	2.5 mL 	5 mL* → 	1 tablet 	_____
36 to 47 pounds (about 16 to 21 kilograms)	4 to 5 years	3.75 mL 	 ← 7.5 mL*	1 ½ tablets 	_____
48 to 59 pounds (about 22 to 26 kilograms)	6 to 8 years	5 mL 	10 mL* → 	2 tablets 	1 tablet 
60 to 71 pounds (about 27 to 32 kilograms)	9 to 10 years	_____	 ← 12.5 mL*	2 ½ tablets 	1 tablet 
72 to 95 pounds (about 33 to 43 kilograms)	11 years	_____	15 mL* → 	3 tablets 	1 ½ tablets 
96 pounds or more (44 kilograms or more)	12 years or older	_____	10 mL + 10 mL = 20 mL* 10 mL* →  and 10 mL* → 	4 tablets 	2 tablets 

***Note:** This dosage is for children's liquid ibuprofen products given by dropper. See the previous column for correct dose-by-weight for infant ibuprofen drops, which are more concentrated.



healthychildren.org

Powered by pediatricians. Trusted by parents.
 from the American Academy of Pediatrics